

An Australian mass casualty incident triage system for the future based on mistakes of the past: The Homebush Triage Standard

Introduction

The combined effects of an aging population, the trend towards day and minimally invasive surgery, plus the increasing cost pressures upon hospitals have significant implications upon the availability of health resources during a mass casualty incident (MCI). There are no universal definitions of what constitutes either a disaster or a MCI within Australia. A 'disaster' is said to have occurred when normal community and organisational arrangements are overwhelmed by an event and extraordinary responses need to be instituted (Emergency Management Australia 1995).

When available medical resources are overwhelmed by casualties, transport and treatment priorities need to be assigned to individuals to ensure limited medical resources are used efficiently. The term triage was transposed from French into the English language during the First World War to describe the process of sorting casualties for treatment priority by the American Army Medical Corps (Rutherford 1989). Casualty triage is the most important medical function during a mass casualty incident (MCI) and accurate triage a major determinant of an individual's survival (Rutherford 1989; Waeckerle 1991; Fryberg *et al* 1988).

This study reviews the evolution of triage, and factors that can potentially interfere with the triage process and compromise the medical response to an MCI. These are then used to synthesize a triage system to provide a common platform so that patient priorities at the incident site can be interfaced with those arising within receiving hospitals.

Historical perspectives MCI triage

The advent of gunpowder and the development of the rifle forced infantry into linear battle formations. As battlefields became larger it became increasingly more difficult to locate wounded soldiers who were left where they fell until

by Dr Antony Nocera, FACEM & Dr Alan Garner, FACEM, NRMA CareFlight Westmead New South Wales

the end of battle. The wounded were then evacuated and treated according to rank including the removal of dead nobles taking priority over wounded common soldiers (Hamby 1967).

Dominique Jean Larrey, Surgeon General to Napoleon's Army of the Rhine, introduced a major revolution in combat casualty care.

Larrey's philosophy was to rescue casualties during battle, with a dedicated corps using purpose built wagons, the ambulance volantes, and rapidly transport them to a central collection point. Here the most seriously wounded would be operated on, without regard to rank or distinction, by either the Surgeon-in-Chief or a competent surgeon under his direction (Richardson 1974). In 1792 Larrey personally lead his ambulance volantes to treat wounded French soldiers in the field and transport them from the front line during the battle against the Austrian Army near Königsberg (Leroy-Dupré 1862).

In 1807 at the Battle of Eylau against the Russian Army, Baron Larrey, now Surgeon-in-Chief to Napoleon's Grand Army, gave treatment based on medical need but with priority to the wounded of Napoleon's Imperial Guard over other wounded French soldiers (Dible 1970). In spite of Larrey's pioneering example, during the American Civil War in 1862, three thousand wounded Union soldiers were left virtually unattended and untreated for three days after the second battle of Bull Run (Adams 1952).

In 1846 British Naval surgeon Dr John Wilson described the principles of MCI triage. Dr Wilson classified combat

injuries into slight, serious and fatal and described a system of treatment priority directed towards the control of life threatening hemorrhage, 'To a serious bleeding everything must of necessity at once give way, and the vessel be secured'. Dr Wilson advocated the treatment of those with fatal injuries be restricted to 'a stimulus, an opiate, a proper easy position' (Wilson 1846).

In the Second World War the procedure of patient triage was regarded as the biggest single factor contributing to survival following abdominal wounds in the US Army (Welch 1947). In the Korean War the application of a four tiered triage system, (immediate, delayed, minimal and expectant) lead to a striking improvement in casualty survival (Hughes 1976). The combination of triage, advanced resuscitation and rapid helicopter evacuation of casualties in the Vietnam War contributed to reducing mortality rates down to 1%, compared to the 4.7% observed during World War Two (Kennedy *et al* 1996).

Goals of MCI triage

The primary objective of military triage was to identify those wounded soldiers who could be treated rapidly and returned to the battlefield (Kennedy *et al* 1996). In civilian practice, the triage process attempts to achieve the greatest good for the greatest number of patients (Emergency Management Australia 1995; Rutherford 1989; Waeckerle 1991; Fryberg *et al* 1988; Burkle 1984). Traditional individual doctor-patient relationships are over-ridden by a collective medical responsibility to the group of casualties (Waeckerle 1991; Burkle 1984; Llewellyn 1992).

In general there is no role for cardio-pulmonary resuscitation during an MCI (Emergency Management Australia 1995) except in cases of lightning strikes involving multiple individuals. Here medical efforts should be directed at those victims in cardio-respiratory arrest, since the majority of other victims will make a

good recovery (Myers et al 1977). Normal triage priorities may be reversed for casualties involved in highly toxic hazardous material exposures where decontamination and treatment priority should be directed at the uninjured and even asymptomatic patients (Kirk et al 1994).

The success of the triage process as a means of minimising preventable deaths during an MCI depends upon being able to rapidly identify those casualties at the extremes of care. Medical resources are diverted from those who will either die, or recover irrespective of the medical care they receive, and concentrated on those critically ill casualties with a reasonable probability of survival (Emergency Management Australia 1995, Waeckerle 1991, Kennedy et al 1996, Burkle 1984).

Problems with MCI Triage

During an MCI, triage is approximately 70% accurate (Burkle 1984) with a tendency to under estimate injury severity. This underscores the need for triage to be viewed as a process of repeated casualty reassessments until the patient receives definitive care. The difficulties in making rapid value judgements based upon relative percentage survival probabilities (Kennedy et al 1996; Hughes 1976; Wardrope et al 1991) adds to the emotional stress upon the individual attempting to perform casualty screening assessments in a hostile environment during an MCI (Spengler 1995).

Triage accuracy is also adversely affected by other factors including, the physiological ability of the young to compensate for hypovolaemia, altered perceptions of pain in high stress situations (Beecher 1946) and neuropsychiatric reactions amongst surviving casualties (Burkle 1996).

MCI Triage considerations for single practitioners in isolated locations

The limited resources and long transportation times dramatically reduce the threshold of what constitutes an MCI in isolated locations and creates unique ethical and practical difficulties in managing incidents. Collective experience from the Korean and Vietnam wars provides some triage guidance in delayed management of penetrating trauma (Moyasenko 1984; Coupland et al 1992).

MCI triage considerations for hospitals

Hospitals must have triage systems to cope with potential incidents in close proximity to their facility where a large number of casualties can present without

Injury	High evacuation and treatment priority	Low evacuation and treatment priority
Penetrating abdominal wounds	Individuals who can access definitive surgical treatment within 6 hours of injury	Survival after 12 hours without operative care
Major vascular injury in an extremity	Individuals who can achieve vessel reconstruction within 10 hours of injury	If vessel reconstruction cannot be achieved within 10 hours direct ligation of the vessel will result in limb loss in 50% of cases.
Penetrating head injuries	Unstable patients with evolving neurological signs	Individuals who are stable, conscious with either no deficit or moderate paresis or hemianopia can survive for 36 hours without neurosurgical care with appropriate fluids, wound care and antibiotics

Table 1: triage considerations for single practitioners in isolated locations

warning before an emergency medical system (EMS) response has been initiated (Caro et al 1973). In addition, large numbers of casualties may be transported directly to the hospital from the incident scene by EMS (Anderson et al 1977) or they may simply overwhelm established EMS field triage & treatment posts, and then move en masse to the nearest hospital (Maningas et al 1997).

The hospital triage process has to be fluid as well as continuous and capable of dealing with incidents where the major casualties are medical rather than surgical (Myers et al 1977; Wardrope et al 1991; Buerk et al 1982). During an MCI hospitals have to integrate casualty triage with the triage of normal daily emergency presentations (De Lorenzo et al 1996). In addition, there may be an increase in normal daily emergency presentations from ischaemic heart disease or exacerbation's of airways disease in response to the occurrence of an incident (Duclos et al 1990; Leor et al 1996).

Special triage situations can arise when hospital resources are either damaged (Schultz et al 1996) or have to be abandoned (De Lorenzo et al 1996, Smith et al 1996) during an incident. Staffing shortages can suddenly arise as health care workers spontaneously evacuate their families in anticipation of a hazard (Smith et al 1981) or cannot reach hospitals due to disruption of transport links. Hospital triage decisions may also have to take into account those outpatients normally dependent on domiciliary medical services who may have to be admitted should an incident temporarily prevent these services from being delivered.

Operational difficulties with MCI triage

Over the years a variety of different triage

systems (Burkle 1984; Coupland et al 1992; Caro et al 1973; Hodgetts et al 1995; Lumley et al 1996; Mac Mahon 1985; Miller 1971; Baskett et al 1988) along with differing triage tag designs to document casualty triage status (Coupland et al 1992; Finch et al 1982; Hodgetts et al 1995; Lumley et al 1996; MacHahon 1985; Baskett et al 1988) have been developed. The crash of a Boeing 737-400 in 1989 at Kegworth in the United Kingdom occurred on the boundary of three different counties whose respective ambulance services used different systems of MCI triage. This incident highlights the avoidable confusion that can arise when responding personnel attempt to use different triage systems and triage tags during an incident (Barton et al 1991; Malone 1990).

Experience has shown that the key operational principle for an efficient disaster response is to ensure staff work as near as practical to their normal daily routines (Pepe et al 1991; Rutherford 1973; Vayer et al 1986). Planning must be based upon what people are likely to do in the stress of an MCI and procedures should be kept simple and practical (Burkle 1984; Mitchell 1986). Accordingly, a triage system must have simple structure and be based on normal daily operating procedures that can also be applied during an MCI.

The Homebush triage taxonomy

There are clear benefits from the standardisation of disaster responses within Australia (Senate Standing Committee on Industry, Science, Technology, Transport, Communications and Infrastructure 1994). A national MCI triage system will mean, in the event of an MCI, both

Note

1. Standards Australia AS-2700 1996 Colour Standard for General Purposes.

hospitals and ambulance services are already using familiar common terminology which will allow effective and efficient communications under stressful circumstances.

A triage system must rapidly screen both children as well as adults (Kennedy *et al* 1996; Klein *et al* 1991), be cost effective, and operable in adverse conditions if they are going to be relevant to single practitioners in isolated areas. The dead and human body parts should be clearly and individually labeled as soon as possible to avoid time being wasted reconfirming death (Rutherford 1989; Burkle 1984) and to prevent the dead being transported to an active treatment area (Faxon 1948).

A simple triage taxonomy with four active treatment levels has been previously used in MCI situations (Hughes 1976; Ammons *et al* 1988; Gans *et al* 1996; Williams *et al* 1974). Increasing the number of categories has not improved the system (Gans *et al* 1996). The Homebush triage taxonomy (table 2) uses these triage priorities as a common core for both prehospital and hospital emergency department operations. Simple mechanisms can expand the four core active treatment groups if required for emergency department quality assurance purposes.

Using standard colors¹ means there will be consistent production standards for triage materials. Providing a common triage language for all healthcare responders eliminates potential communication problems associated with using different terminology. This will facilitate the integration of military medical services in the event they were deployed to assist the civilian response of a large-scale MCI within a State. A single common triage

Classification	Priority	Priority code	Colour	Colour number ¹
Immediate	Top	A (Alpha)	Red (Homebush Red)	R 22
Urgent	High	B (Bravo)	Yellow (Homebush Gold)	Y 26
NOT Urgent	Low	C (Charlie)	Green (Homebush Green)	G 27
Dying	Terminal Care only	D (Delta)	White	N 14
Dead	Not Applicable	E (Echo)	Black	N 61

Table 2: The Homebush triage taxonomy

system reduces the problems that military personnel would also face trying to interface with different civilian medical services especially with a large number of medical evacuations across State borders, where different systems of MCI triage are used.

The use of phonetic triage priority codes instead of numerical codes takes into account the problems with radio voice transmission. Numbers are reserved to either stratify patient priorities within a particular triage category, or to quantify the number of casualties within a particular triage category.

In an overwhelming situation there will be patients for whom the difficult decision not to treat must be made (Parke *et al* 1992). However the decision on what constitutes a non-survivable injury is a balance between the magnitude of the incident, an individual casualty's relative probability of survival, and the capacity of available medical resources at different points in the casualty evacuation chain (Waeckerle 1991; Llewellyn 1992).

A specific triage category for dying patients provides clear management

directions for those patients assessed as being beyond help either at the incident site (Fryberg 1988), the casualty collection point or emergency department (Rutherford 1989; Williams *et al* 1974; Sharpe 1985; Artuson 1981; Das 1983; Seletz 1990) or on the operating table (Burkle *et al* 1994). The introduction of this classification into daily emergency department operations identifies those patients with advance medical treatment directives and directs appropriate care to patients with terminal chronic illnesses.

Homebush Triage methodology

The Simple Triage and Rapid Treatment (START) and Secondary Assessment of Victim Endpoint (SAVE) (Benson *et al* 1996) attempt to apply the principles of evidence based medicine to disaster triage. START triage has been used successfully at several MCIs within the United States. These include the 1995 Oklahoma City Bombing, the 1992 Bombing of the New York World Trade Center, Hurricane Andrew, and the 1989 Northridge earthquake (Personal communication Dr Carl Schultz).




Table 3: The Simple Triage And Rapid Treatment (START) Triage scheme (modified). Combining START with the Homebush triage taxonomy allows a simple triage decision tree to be developed. Reprinted with the permission of Prehospital and Disaster Medicine.

The simplicity of START (table 3) allows it to be performed rapidly as a quick screening tool and can be easily remembered as:

- anyone who does not breathe with simple airway maneuvers is dead
- anyone who can walk is assigned a not urgent triage priority
- anyone who cannot walk but can obey commands, with both a radial pulse being present and a respiratory rate less than 30 breaths per minute, is assigned an urgent triage priority
- anyone else has an immediate triage priority

SAVE guidelines look at a number of parameters (table 4) which are designed to answer two key triage questions at a major incident site (Benson *et al* 1996):

- What is the victim's prognosis if minimal treatment is provided?
- What is the victim's prognosis with the treatment resources available at the area medical center?

There has never been a situation to date that has required the implementation of both START and SAVE triage criteria.

Paediatric triage

The basic principles of triage remain the same for children as they are for adults (Holbrook 1991). The START methodology will tend to overtriage children. This is acceptable given the higher probability of children surviving head injury (Luerksen *et al* 1988) and multiorgan system failure compared with adults (Wilkinson *et al* 1986), along with the fact that most blunt abdominal trauma is managed conservatively in children compared with adults (Powell *et al* 1987). The initial Glasgow Coma Score following head injury in children does not reliably predict outcome unless there is associated hypoxia and hypotension present (Lieh-Lai *et al* 1992).

Triage documentation

Triaging patients into geographic areas has been raised as an alternative to the use of triage tags (Rutherford 1989; Vayer *et al* 1986; Vukmir *et al* 1991; Kerns *et al* 1990). Geographic triage provides a major time saving in triage documentation especially when there is a large influx of patients. (Waeckerle 1991; Kennedy *et al* 1996; Vayer *et al* 1986; Angus *et al* 1993). In a series of six major air accidents within the USA the largest incident, involving 297 people with 59 critically injured and 124 less severely injured, had the shortest prehospital time using geographic triage instead of triage tags, combined with efficient ground and rotary wing transport systems (Anderson 1995).

SAVE Guidelines

- Mangled Extremity Severity Score (MESS) (Johansen *et al* 1990) to assess crush injury to extremities
- Glasgow Coma Score less than eight in adults with significant head injury.
- Abdominal trauma with refractory hypotension
- Chest trauma with abnormal vital signs
- Spinal trauma
- Burns with < 50% probability of survival or adults over 60 years of age with an inhalational injury.
- Adults with pre-existing diseases
- Non traumatic emergencies
- Special triage categories such as healthcare workers with minor injuries who with simple treatment may be able to assist in the medical response

Table 4: Secondary Assessment of Victim Endpoint (SAVE) Guidelines

Triage is generally carried out once the casualties have been taken to a casualty collection point (Burkle 1984; Vayer *et al* 1986; Orr *et al* 1983). Triage flags provide the first ambulance on scene at an MCI with the capability to lay the foundation for the site medical response irrespective of the number of casualties. Geographic triage may reflect the normal disposition of trauma patients at an incident (Vukmir *et al* 1991) that can assist single practitioners with limited site resources to triage casualties efficiently.

The casualty profile following an MCI typically has 6 to 25% of patients requiring medical or surgical treatment within 12 hours to prevent loss of life or severe morbidity (Anderson 1995; Sklar 1987). The bulk of the casualty load consists of patients with non-urgent injuries who have little to gain from immediate medical care. Using expensive triage tags to identify them or label dead bodies is an inappropriate use of resources especially in a large scale MCI (Rutherford 1989; Waeckerle 1991; Angus *et al* 1993). In 1974 a Turkish DC10 crashed into a forest at Ermenoville, France killing 345 persons. Nearly 20,000 fragments of human tissue were produced from the impact with the remains of 188 victims subsequently positively identified (Personal communication Mr. Peter J. Stuart).

Triage procedures should avoid unnecessarily complicating the subsequent investigation of the incident. Labelling human remains with numbered chemically resistant tags helps to document the location of human body parts and their relationship to objects such as motor vehicles at the scene. This facilitates their systematic removal from the site for subsequent forensic examination and can play an important role in victim identification.

Conclusion

Australia has the opportunity to build upon past experience and develop a nationally integrated system of casualty triage. Appropriate preplanning can mitigate some of the problems that complicate the triage process, but those involved in the medical response to an MCI must have a common language and understanding of triage issues to remove existing fundamental barriers to good communications.

References

- Adams G. W. 1952, 'Doctors in blue', *The medical history of the Union Army in the Civil War*, Henry Schuman, New York.
- Ammons M. A., Moore E. E., Pons P. T., & Moore F.A *et al* 1988, 'The role of a regional trauma system in the management of a mass disaster: an analysis of the Keystone, Colorado, chairlift accident', *J Trauma*, 28, pp. 1468-1471.
- Anderson G.V. & Feliciano D. V. 1977, 'The Centennial Olympic Park bombing: Grady's Response', *J Med Assoc Ga*, 86, pp. 42-46.
- Anderson P. B. 1995, 'A comparative analysis of the emergency medical services and rescue responses to eight airliner crashes in the United States, 1987-1991', *Prehospital & Disaster Medicine*, 10, pp. 142-153.
- Angus D. C. & Kvetan V. 1993, 'Organization and management of critical care systems in unconventional situations', *Crit Care Clin*, 9, pp. 521-542.
- Arturson G. 1981, 'The Los Alfaques disaster: a boiling-liquid, expanding-vapour explosion', *Burns*, 7, pp. 233-251.
- Barton D, Bodiwala G.G. 1991, 'Assessment of a triage labels system during a major incident exercise', *Prehospital and Disaster medicine*, 6, pp. 473-476.
- Baskett P. & Wells R. (eds.) 1988,

Medicine for disasters, Wright, London.

Beecher H. K. 1946, 'Pain in men wounded in battle', *Ann Surg*, 123, pp. 96–105.

Benson M., Koenig K. L. & Schultz C. H. 1996, 'Disaster triage: START, then SAVE—A new method of dynamic triage for Victims of a Catastrophic Earthquake', *Prehospital and Disaster Medicine*, 11, pp. 117–24.

Buerk C. A., Batdorf J. W., Cammack K. V., & Ravenbolt O. 1982, 'The MGM Grand Hotel fire. Lessons learned from a major disaster', *Arch Surg*, 117, pp. 641–644.

Burkle F. M. 1996, 'Acute-Phase mental health consequences of disasters: Implications for triage & emergency medical services', *Ann Emerg Med*, 28, pp. 119–128.

Burkle F. M. (senior editor) 1994, *Disaster medicine. Application for the immediate management and triage of civilian and military disaster victims*, Medical Examination Publishing Company, New York.

Burkle F. M., Orebaugh S., & Barendse B. R. 1994, 'Emergency medicine in the Persian Gulf War—Part 1: Preparations for triage and combat casualty care', *Ann Emerg Med*, 23, pp. 742–747.

Caro D. & Irving M. 1973, 'The Old Bailey bomb explosion', *Lancet*, 1, pp. 1433–1435.

Coupland R. M., Parker P.J. & Gray R. C. 1992, 'Triage of war wounded: the experience of the International Committee of the Red Cross', *Injury*, 23, pp. 507–510.

Das R. A. P. 1983, '1981 circus fire disaster in Bangalore, India: Causes, management of burn patients and possible presentation', *Burns*, 10, pp. 17–29.

De Lorenzo R. A. & Augustine J. J. 1996, 'Lessons in emergency evacuation from the Maimsburg train derailment', *Prehospital & disaster medicine*, 11, pp. 270–275.

Dible J. H. 197, *Napoleon's Surgeon*, William Heinemann Medical Books Ltd, London.

Duclos P., Sanderson L.M. & Lipsett M. 1990, 'The 1987 forest fire disaster in California: assessment of emergency room visits', *Arch of Environ Health*, 45, pp. 53–58.

Emergency Management Australia 1995, *Australian Emergency Manual. Disaster Medicine*, Commonwealth Department of Human Services & Health, Canberra.

Faxon N. W. 1948, 'The problems of the hospital administration—The Coconut Grove disaster', *Ann Surg*, 117, pp. 803–808.

Finch P.M. & Hamilton T. 1982, 'Early documentation of disaster victims', *Anaesthesia*, 37, pp. 1185–1189.

Frykberg E. R. & Tepas J. J. 1988, 'Terrorist bombings Lessons learned from Belfast to Beirut', *Ann Surg*, 208, pp. 569–576.

Gans L. & Kennedy T. 1996, 'Management

of unique clinical entities in disaster medicine', *Emerg Med Clin North Am*, 14, pp. 301–326.

Hamby W. B. 1967, *Ambrosio Paré. Surgeon of the Renaissance*, William H. Green Inc., St Louis.

Hodgetts T. J. & Mackway-Jones K. (eds.) 1995, *Major incident medical management & support. The practical approach*, BMJ Publishing Group, Plymouth.

Holbrock P. R. 1991, 'Pediatric disaster medicine'. *Crit Care Clin*, 7 pp. 463–470.

Hughes J. H. 1976, 'Community medicine. Triage—A new look at an old French concept', *Postgrad Med*, 60, pp. 223–227.

Johansen K., Daines M., Howey T., Helfet D. & Hansen S. T. 1990, 'Objective criteria accurately predict amputation following lower extremity trauma', *J Trauma*, 30, pp. 568–572.

Kennedy K., Aghababian R. V., Gans L. & Lewis C. P. 1996, 'Triage: Techniques and applications in decision making', *Ann Emerg Med*, 28, pp. 136–144.

Kerns D. E. & Anderson P. B. 1990, 'EMS response to a major aircraft incident: Sioux City, Iowa', *Prehospital and Disaster Medicine*, 5, pp. 159–166.

Kirk M. A., Cisek J. & Rose S. R. 1994, 'Emergency Department response to hazardous material incidents', *Emerg Med Clin North Am*, 12, pp. 461–481.

Klein J. S. & Weigelt J. A. 1991, 'Disaster management lessons learned', *Surg Clin North Am*, 71, pp. 257–266.

Leor J., Poole W.K. & Kloner R. A. 1996, 'Sudden cardiac death triggered by an earthquake', *N Eng J Med*, 334, pp. 413–419.

Memoir of Baron Larrey, Surgeon-in-Chief of the Grande Armée. From the French (of L.A.H. Leroy-Dupré). Second edition. Henry Renshaw, London 1862.

Lieh-Lai M.W., Theodorou A.A., Sarnaik A.P., Meert K.L., Moylan P.M. & Canady A.I. 1992, 'Limitations of the Glasgow Coma Scale in predicting outcome in children with traumatic brain injury', *J Pediatr*, 120, pp. 195–199.

Llewellyn C. H. 1992, 'Triage: in austere environments and echeloned medical systems', *World J Surg*, 16, pp. 904–909.

Luerssen T. G., Klauber M.R. & Marshall L. F. 1998, 'Outcome from head injury related to patient's age. A longitudinal prospective study of adult and pediatric head injury', *J Neurosurg*, 68, pp. 409–416.

Lumley J. S. P., Ryan J. M., Baxter P. J. & Kirby N. 1996, *Handbook of the Medical Care of Catastrophes*, Royal Society of Medicine Press, London.

MacMahon A. G. 1985, 'Sorting out triage in urban disasters', *SAMJ*, 67, pp. 555–556.

Malone W.D. 1990, 'Lessons to be learned from the major disaster following the civil

airliner crash at Kegworth in January 1989', *Injury*, 21, pp. 49–52.

Maningas P.A., Robinson M., Mallonee S. 'The EMS response to the Oklahoma City Bombing', *Prehospital and Disaster Medicine*, 1997, 12, pp. 9–14.

Miller P. J. 1971, 'The management of major incidents', *Injury*, 2, pp. 68–181.

Mitchell G. W. 1986, 'The triage process', *Topics in Emergency Medicine*, 7, pp. 34–45.

Moysaenko V. 1984, 'Multiple trauma Chapter 19 of: Burkle FM Senior editor. Disaster medicine', *Application for the immediate management and triage of civilian and military disaster victims*, Medical Examination Publishing Company, New York.

Myers G.J., Colgan M. T. & VanDyke D. H. 1977, 'Lightning-strike disaster among children', *JAMA*, 238, pp. 1045–1046.

Orr S. M. & Robinson W. A. 1983, 'The Hyatt Regency skywalk collapse: an EMS-based disaster response', *Ann Emerg Med*, 12, pp. 601–605.

Parke T.R., Haddock G., Steedman D. J., Pollock A. J. & Little K. 1992, 'Response to the Kurdish refugee crisis by the Edinburgh MEDIC 1 team', *BMJ*, 304, pp. 695–697.

Pepe P. E. & Kvetan V. 1991, 'Field management and critical care in mass disasters', *Crit Care Clin*, 7, pp. 401–420.

Powell R.W., Green J. B., Ochsner M. G., Barttelbort S. W., Shackford S. R. & Sise M. J. 1987, 'Peritoneal lavage in pediatric patients sustaining blunt abdominal trauma: a reappraisal', *J Trauma*, 27, pp. 6–9.

Richardson R. G. 1974, *Larrey: Surgeon to Napoleon's Imperial Guard*, John Murray, London.

Rutherford W. H. 1973, 'Experience in the accident and emergency department of the Royal Victoria Hospital with patients from civil disturbances in Belfast 1969–1972, with a review of disasters in the United Kingdom 1951–1971', *Injury*, 4, pp. 189–199.

Rutherford W. 1989, 'Triage for simple compensated disasters', *Journal British Association for Immediate Care*, 12, pp. 62–67.

Schultz C. H., Koenig K. L. & Noji E. K. 1996, 'A medical disaster response to reduce immediate mortality after an earthquake', *N Eng J Med*, 334, pp. 438–444.

Seletz J. M. 1990, 'Flugtag-88 (Ramstein Air Show Disaster): an Army response to a MASCAL', *Mil Med*, 155, pp. 152–155.

Senate Standing Committee on Industry, Science, Technology, Transport, Communications and Infrastructure. Chair Senator BK Childs, 1994, *Disaster Management*, The Parliament of the Commonwealth of Australia.

Sharpe, D. T., Roberts A. H., Barclay T. L., Dickson W. A., Settle J. A. D., Crockett D. J. & Mossad M. G. 1985, 'Treatment of burns casualties after fire at Bradford City football ground', *BMJ*, 291, pp. 945-948.

Sklar D. P. 1987, 'Casualty patterns in disasters', *Journal World Association Emergency and Disaster Medicine*, 3, pp. 49-51.

Smith J. S. & Fisher J. H. 1981, 'Three Mile Island. The silent disaster', *JAMA*, 245, pp. 1656-1659.

Spengler C. 1995, 'The Oklahoma City bombing: a personal account' *J Child Neurol*, 10, pp. 392-398.

Vayer J. S., Ten Eyck R. P. & Cowan M. L. 1986, 'New concepts in triage', *Ann Emerg Med*, 15, pp. 27-30.

Vukmir R. B. & Paris P. M. 1991, 'The Three Rivers Regatta accident: an EMS

perspective', *Am J Emerg Med*, 9, pp. 64-71.

Waeckerle J. F. 1991, 'Disaster planning & response', *N Eng J Med*, 324, pp. 815-821.

Wardrope J., Ryan E., Clark G., Venables G., Crosby A. C. & Redgrave P. 1991, 'The Hillsborough tragedy', *BMJ*, 303, pp. 1381-1385.

Welch C. E. 1947, 'War wounds of the abdomen', *N Eng J Med*, 1, pp. 156-162.

Williams R. W., Burns G. P., Andersen M. N., Reading G. P. & Border J. R. *et al* 1974, 'Mass casualties in a maximum security institution', *Ann Surg*, 179, pp. 592-597.

Wilson J. (D.) 1846, *Outlines of Naval surgery*, Maclachlan, Stewart and Co, Edinburgh.

Wilkinson J. D., Pollack M. M., Ruttiman U. E., Glass N. L. & Yeh T. S. 1986, 'Outcome of pediatric patients with multiple organ system failure', *Crit Care Clin*, 14, pp. 271-274.

Acknowledgement

The authors would like to acknowledge the assistance of Ms Anne Newton in the formulation of the manuscript.

Declaration

This paper has been prepared without any funding from any organization or institution. The original manuscript was exclusively submitted to the Australian and New Zealand Journal of Surgery as an original work by the authors who retain the copyright over the manuscript.

Author's contact details

Dr Antony Nocera, FACEM & Dr Alan Garner, FACEM
NRMA Careflight
PO Box 159
Westmead NSW 2145

Dr Antony Nocera
Unit 16, 68 Bradley's Head Road
Mosman NSW 2088
tonynoc@ozemail.com.au
Tel. (Mobile): 0414-959599

EMA Safer Communities Awards

EMA

Safer Communities Awards

From flood recovery processes to innovations in firefighting, the entries in this year's Emergency Management Australia (EMA) Safer Communities Awards have shown excellence in many areas of emergency management.

More than 75 entries were received from across Australia. They were submitted by State Government, Local Government, private sector and volunteer organisations, and covered both pre and post-disaster emergency management.

The innovation and leadership exhibited in the entries demonstrated real achievement in helping communities prepare for, as well as recover from, disasters such as flooding, cyclones or bushfires.

Following the selection of 26 winners at a State and Territory level, the judging panel chose the following eight National winners, with five commendations made to other entries. The winners were presented with their Awards at a ceremony hosted by the Minister for Defence, Mr John Moore MP, at Parliament House, Canberra, on 28 June 2000.

Department of Human Services, Gippsland (VIC)

Post-disaster category
Federal/State Government Stream

For its pro-active response in resourcing, implementing and monitoring the immediate welfare and recovery activities following the severe flooding of the East Gippsland Shire in 1998.

Wollongong City Council (NSW)

Post-disaster category
Local Government Stream

For the work of their geo-technical team which had the task of assessing landslide risks following flash flooding in the area. The team's knowledge and expertise saved valuable resources by evacuating only those who absolutely had to be evacuated.